


CLIMATE CHANGE PROJECTIONS FOR INDIVIDUAL TREE SPECIES SOUTHERN AND COASTAL NEW ENGLAND



This region's forests will be affected by a changing climate and other stressors during this century. A team of managers and researchers created an assessment that describes the vulnerability of forests in the region ([Janowiak et al. 2018](#)). This report includes information on observed and future climate trends, and also summarizes key vulnerabilities

for forested natural communities. The Landscape Change Research Group recently updated the Climate Change Tree Atlas, and this handout summarizes that information. Full Tree Atlas results are available online at www.fs.fed.us/nrs/atlas/. Two climate scenarios are presented to "bracket" a range of possible futures. These future climate projections (2070 to 2099) provide information about how individual tree species may respond to a changing climate. Results for "low" and "high" emissions scenarios can be compared on the reverse side of this handout.

The updated Tree Atlas presents additional information helpful to interpret tree species changes:

- **Suitable habitat** - calculated based on 39 variables that explain where optimum conditions exist for a species, including soils, landforms, and climate variables.
- **Adaptability** - based on life-history traits that might increase or decrease tolerance of expected changes, such as the ability to withstand different forms of disturbance.
- **Capability** - a rating of the species' ability to cope or persist with climate change in this region based on suitable habitat change (statistical modeling), adaptability (literature review and expert opinion), and abundance (FIA data). The capability rating is modified by abundance information; ratings are downgraded for rare species and upgraded for abundant species.
- **Migration Potential Model** - when combined with habitat suitability, an estimate of a species' colonization likelihood for new habitats. This rating can be helpful for assisted migration or focused management (see the table section: "New Habitat with Migration Potential").

Remember that models are just tools, and they're not perfect. Model projections can't account for all factors that influence future species success. If a species is rare or confined to a small area, model results may be less reliable. These factors, and others, could cause a particular species to perform better or worse than a model projects. Human choices will also continue to influence forest distribution, especially for tree species that are projected to increase. Planting programs may assist the movement of future-adapted species, but this will depend on management decisions. Despite these limits, models provide useful information about future expectations. It's perhaps best to think of these projections as indicators of possibility and potential change.

SOURCE: This handout summarizes the full model results for the Southern and Coastal New England region, available at www.fs.fed.us/nrs/atlas/combined/resources/summaries. More information on vulnerability and adaptation in the New England region can be found at www.forestadaptation.org/new-england. A full description of the models and variables are provided in Iverson et al. 2019 (www.nrs.fs.fed.us/pubs/57857) and Peters et al. 2019 (www.nrs.fs.fed.us/pubs/58353).

CLIMATE CHANGE CAPABILITY

POOR CAPABILITY

American basswood	Paper birch
Balsam fir	Pitch pine
Balsam poplar	Red pine
Black ash	Red spruce
Black spruce	Slippery elm
Black walnut	Striped maple
Bur oak	Swamp white oak
Eastern white pine	Sweet birch
Hackberry	Tamarack (native)
Northern pin oak	White spruce
Northern white-cedar	

FAIR CAPABILITY

Black locust	White ash
Quaking aspen	

GOOD CAPABILITY

American beech	Post oak
American holly	Red maple
Black cherry	Sassafras
Black oak	Scarlet oak
Blackgum	Shagbark hickory
Chestnut oak	Sugar maple
Eastern redcedar	Sweetgum
Mockernut hickory	White oak
Northern red oak	Yellow-poplar
Pignut hickory	

MIXED RESULTS

American elm	Gray birch
American hornbeam	Green ash
Bigtooth aspen	Ironwood
Eastern hemlock	Silver maple
Flowering dogwood	Yellow birch

NEW HABITAT WITH MIGRATION POTENTIAL

Chinkapin oak	Sweetbay
Loblolly pine	Virginia pine
Shortleaf pine	Water hickory
Southern red oak	



ADAPTABILITY: Life-history factors, such as the ability to respond favorably to disturbance, that are not included in the Tree Atlas model and may make a species more or less able to adapt to future stressors.

- + **HIGH** *Species may perform better than modeled*
- **MEDIUM**
- **LOW** *Species may perform worse than modeled*

HABITAT CHANGE: Projected change in suitable habitat between current and potential future conditions.

- ▲ **INCREASE** *Projected increase of >20% by 2100*
- **NO CHANGE** *Projected change of <20% by 2100*
- ▼ **DECREASE** *Projected decrease of >20% by 2100*
- ★ **NEW HABITAT** *Tree Atlas projects new habitat for species not currently present*

ABUNDANCE: Based on Forest Inventory Analysis (FIA) summed Importance Value data, calibrated to a standard geographic area.

- + **ABUNDANT**
- **COMMON**
- **RARE**

CAPABILITY: An overall rating that describes a species' ability to cope or persist with climate change based on suitable habitat change class (statistical modeling), adaptability (literature review and expert opinion), and abundance within this region.

- ▲ **GOOD** *Increasing suitable habitat, medium or high adaptability, and common or abundant*
- **FAIR** *Mixed combinations, such as a rare species with increasing suitable habitat and medium adaptability*
- ▼ **POOR** *Decreasing suitable habitat, medium or low adaptability, and uncommon or rare*

SPECIES	LOW CLIMATE CHANGE (RCP 4.5)				HIGH CLIMATE CHANGE (RCP 8.5)			
	ADAPT	ABUN	HABITAT CHANGE	CAPABILITY	ADAPT	ABUN	HABITAT CHANGE	CAPABILITY
American basswood	•	-	●	▼	●	▼	●	▼
American beech	•	•	▲	▲	▲	▲	▲	▲
American elm	•	•	▼	▼	●	○	▼	▼
American holly	•	-	▲	▲	▲	▲	▲	▲
American hornbeam*	•	-	●	▼	▲	▲	▲	▲
American mountain-ash*	-	-	▼	▼	▼	▼	▼	▼
Atlantic white-cedar*	-	-	●	▼	●	▼	●	▼
Bald cypress	•	-	★		★			
Balsam fir	-	•	▼	▼	▼	▼	▼	▼
Balsam poplar	•	-	▼	▼	▼	▼	▼	▼
Bigtooth aspen	•	•	▲	▲	●	○	▼	▼
Bitternut hickory*	+	-	●	○	▲	▲	▲	▲
Black ash	-	-	▼	▼	▼	▼	▼	▼
Black cherry	-	•	▲	▲	▲	▲	▲	▲
Black locust*	•	-	▲	○	▲	○	▲	○
Black oak	•	•	▲	▲	▲	▲	▲	▲
Black spruce	•	-	▼	▼	▼	▼	▼	▼
Black walnut*	•	-	▼	▼	●	▼	▼	▼
Blackgum	+	-	▲	▲	▲	▲	▲	▲
Boxelder*	+	-	●	○	●	○	●	○
Bur oak	+	-	▼	▼	▼	▼	▼	▼
Chestnut oak	+	-	▲	▲	▲	▲	▲	▲
Chinkapin oak	•	-			★		★	
Eastern hemlock	-	+	▼	○	▼	▼	▼	▼
Eastern redcedar	•	•	▲	▲	▲	▲	▲	▲
Eastern white pine	-	+	▼	▼	▼	▼	▼	▼
Flowering dogwood	•	-	●	▼	▲	○	▲	○
Gray birch*	•	-	●	▼	▲	○	▲	○
Green ash*	•	-	▲	○	▲	▲	▲	▲
Hackberry	+	-	▼	▼	▼	▼	▼	▼
Honeylocust*	+	-	▼	▼	▼	▼	▼	▼
Ironwood*	+	-	●	○	▲	▲	▲	▲
Loblolly pine	•	-	★		★			
Longleaf pine	•	-	★		★			
Mockernut hickory	+	-	▲	▲	▲	▲	▲	▲
Northern pin oak	+	-	▼	▼	▼	▼	▼	▼
Northern red oak	+	+	●	▲	▼	▲	▼	▲
Northern white-cedar	•	-	▼	▼	▼	▼	▼	▼
Paper birch	•	•	▼	▼	▼	▼	▼	▼
Pignut hickory	•	•	▲	▲	▲	▲	▲	▲
Pin cherry*	•	-	▼	▼	▼	▼	▼	▼
Pin oak*	-	-	●	▼	▲	▲	○	○
Pitch pine	•	•	▼	▼	▼	▼	▼	▼
Post oak	+	-	▲	▲	▲	▲	▲	▲
Quaking aspen	•	•	●	○	●	○	●	○
Red maple	+	+	▼	▲	▼	▲	▼	▲
Red pine	-	-	▼	▼	▼	▼	▼	▼
Red spruce	-	-	▼	▼	▼	▼	▼	▼
Sassafras*	•	-	▲	▲	▲	▲	▲	▲
Scarlet oak	•	•	▲	▲	▲	▲	▲	▲
Shagbark hickory	•	•	▲	▲	▲	▲	▲	▲
Shortleaf pine	•	-	★		★			
Silver maple*	+	-	▼	▼	●	○	▼	○
Slippery elm*	•	-	▼	▼	▼	▼	▼	▼
Southern red oak	+	-	★		★			
Striped maple	•	-	▼	▼	▼	▼	▼	▼
Sugar maple	+	•	▲	▲	▲	▲	▲	▲
Swamp chestnut oak*	•	-	▼	▼	▲	▲	▲	▲
Swamp tupelo	-	-	★		★			
Swamp white oak*	•	-	●	▼	●	▼	▼	▼
Sweet birch	-	•	●	▼	▼	▼	▼	▼
Sweetbay	•	-	★		★			
Sweetgum	•	-	▲	▲	▲	▲	▲	▲
Sycamore*	•	-	●	▼	▲	○	▲	○
Tamarack (native)	-	-	▼	▼	▼	▼	▼	▼
Virginia pine	•	-	★		★			
Water hickory	•	-			★			
White ash	-	•	▲	○	▲	○	▲	○
White oak	+	•	▲	▲	▲	▲	▲	▲
White spruce	•	-	▼	▼	▼	▼	▼	▼
Yellow birch	•	•	●	○	▲	▲	▲	▲
Yellow-poplar	+	-	▲	▲	▲	▲	▲	▲

*Species with low model reliability based on five statistical metrics of the habitat models that affect change class. See maps and tables for more information (www.fs.fed.us/nrs/atlas/combined/resources/summaries).