


CLIMATE CHANGE PROJECTIONS FOR INDIVIDUAL TREE SPECIES NEW ENGLAND AND NORTHERN NEW YORK



This region's forests will be affected by a changing climate and other stressors during this century. A team of managers and researchers created an assessment that describes the vulnerability of forests in the region ([Janowiak et al. 2018](#)). This report includes information on observed and future climate trends, and also summarizes key vulnerabilities

for forested natural communities. The Landscape Change Research Group recently updated the Climate Change Tree Atlas, and this handout summarizes that information. Full Tree Atlas results are available online at www.fs.fed.us/nrs/atlas/. Two climate scenarios are presented to "bracket" a range of possible futures. These future climate projections (2070 to 2099) provide information about how individual tree species may respond to a changing climate. Results for "low" and "high" emissions scenarios can be compared on the reverse side of this handout.

The updated Tree Atlas presents additional information helpful to interpret tree species changes:

- **Suitable habitat** - calculated based on 39 variables that explain where optimum conditions exist for a species, including soils, landforms, and climate variables.
- **Adaptability** - based on life-history traits that might increase or decrease tolerance of expected changes, such as the ability to withstand different forms of disturbance.
- **Capability** - a rating of the species' ability to cope or persist with climate change in this region based on suitable habitat change (statistical modeling), adaptability (literature review and expert opinion), and abundance (FIA data). The capability rating is modified by abundance information; ratings are downgraded for rare species and upgraded for abundant species.
- **Migration Potential Model** - when combined with habitat suitability, an estimate of a species' colonization likelihood for new habitats. This rating can be helpful for assisted migration or focused management (see the table section: "New Habitat with Migration Potential").

Remember that models are just tools, and they're not perfect. Model projections can't account for all factors that influence future species success. If a species is rare or confined to a small area, model results may be less reliable. These factors, and others, could cause a particular species to perform better or worse than a model projects. Human choices will also continue to influence forest distribution, especially for tree species that are projected to increase. Planting programs may assist the movement of future-adapted species, but this will depend on management decisions. Despite these limits, models provide useful information about future expectations. It's perhaps best to think of these projections as indicators of possibility and potential change.

SOURCE: This handout summarizes the full model results for the New England and Northern New York region, available at www.fs.fed.us/nrs/atlas/combined/resources/summaries. More information on vulnerability and adaptation in the New England region can be found at www.forestadaptation.org/new-england. A full description of the models and variables are provided in Iverson et al. 2019 (www.nrs.fs.fed.us/pubs/57857) and Peters et al. 2019 (www.nrs.fs.fed.us/pubs/58353).

CLIMATE CHANGE CAPABILITY

POOR CAPABILITY

Balsam fir	Pin cherry
Balsam poplar	Pitch pine
Black ash	Red pine
Black spruce	Serviceberry
Bur oak	Striped maple
Northern white-cedar	Tamarack (native)

FAIR CAPABILITY

Bitternut hickory	Red spruce
Boxelder	Shagbark hickory
Eastern hemlock	Silver maple
Eastern white pine	Sweet birch
Gray birch	White spruce
Jack pine	Yellow birch

GOOD CAPABILITY


American basswood	Ironwood
American beech	Mockernut hickory
American holly	Northern pin oak
Bigtooth aspen	Nothern red oak
Black cherry	Post oak
Black locust	Quaking aspen
Black oak	Red maple
Blackgum	Sugar maple
Chestnut oak	Sweetgum
Eastern redcedar	White oak
Hackberry	Yellow-poplar

MIXED RESULTS

American elm	Pignut hickory
Paper birch	Scarlet oak

NEW HABITAT WITH MIGRATION POTENTIAL

Chinkapin oak	Shortleaf pine
Common persimmon	Southern red oak
Cucumbertree	Sweetbay
Eastern redbud	Virginia pine
Loblolly pine	Willow oak
Osage-orange	



ADAPTABILITY: Life-history factors, such as the ability to respond favorably to disturbance, that are not included in the Tree Atlas model and may make a species more or less able to adapt to future stressors.

- + **HIGH** *Species may perform better than modeled*
- **MEDIUM**
- **LOW** *Species may perform worse than modeled*

HABITAT CHANGE: Projected change in suitable habitat between current and potential future conditions.

- ▲ **INCREASE** *Projected increase of >20% by 2100*
- **NO CHANGE** *Projected change of <20% by 2100*
- ▼ **DECREASE** *Projected decrease of >20% by 2100*
- ★ **NEW HABITAT** *Tree Atlas projects new habitat for species not currently present*

ABUNDANCE: Based on Forest Inventory Analysis (FIA) summed Importance Value data, calibrated to a standard geographic area.

- + **ABUNDANT**
- **COMMON**
- **RARE**

CAPABILITY: An overall rating that describes a species' ability to cope or persist with climate change based on suitable habitat change class (statistical modeling), adaptability (literature review and expert opinion), and abundance within this region.

- ▲ **GOOD** *Increasing suitable habitat, medium or high adaptability, and common or abundant*
- **FAIR** *Mixed combinations, such as a rare species with increasing suitable habitat and medium adaptability.*
- ▼ **POOR** *Decreasing suitable habitat, medium or low adaptability, and uncommon or rare*

SPECIES	ADAPT		LOW CLIMATE CHANGE (RCP 4.5)		HIGH CLIMATE CHANGE (RCP 8.5)	
	ABUN	CHANG	HABITAT	CAPABILITY	HABITAT	CAPABILITY
American basswood	•	-	▲	△	▲	△
American beech	•	+	●	△	●	△
American elm	•	•	●	○	▲	△
American holly	•	-	▲	△	▲	△
American hornbeam*	•	-	●	▼	●	▼
Atlantic white-cedar*	-	-	●	▼	●	▼
Bald cypress	•	-	★		★	
Balsam fir	-	+	▼	▼	▼	▼
Balsam poplar	•	-	▼	▼	▼	▼
Bigtooth aspen	•	•	▲	△	▲	△
Bitternut hickory*	+	-	●	○	●	○
Black ash	-	-	▲	▼	▲	▼
Black cherry	-	•	▲	△	▲	△
Black locust*	•	-	▲	△	▲	△
Black oak	•	•	▲	△	▲	△
Black spruce	•	•	▼	▼	▼	▼
Black walnut*	•	-	▲	○	▲	△
Blackgum	+	-	▲	△	▲	△
Boxelder*	+	-	●	○	●	○
Bur oak	+	-	▼	▼	▼	▼
Chestnut oak	+	-	▲	△	▲	△
Chinkapin oak	•	-	★		★	
Common persimmon*	+	-	★		★	
Cucumber tree*	•	-	★		★	
Eastern hemlock	-	+	●	○	●	○
Eastern redbud*	•	-	★		★	
Eastern redcedar	•	-	▲	△	▲	△
Eastern white pine	-	+	●	○	●	○
Flowering dogwood	•	-	●	▼	▲	△
Gray birch*	•	•	●	○	●	○
Green ash*	•	-	●	▼	▲	○
Hackberry	+	-	▲	△	▲	△
Ironwood*	+	•	▲	△	▲	△
Jack pine	+	-	●	○	●	○
Loblolly pine	•	-	★		★	
Longleaf pine	•	-	★		★	

SPECIES	ADAPT		LOW CLIMATE CHANGE (RCP 4.5)		HIGH CLIMATE CHANGE (RCP 8.5)	
	ABUN	CHANG	HABITAT	CAPABILITY	HABITAT	CAPABILITY
Mockernut hickory	+	-	▲	△	▲	△
Mountain maple*	+	-	▼	▼	▼	▼
Northern pin oak	+	-	▲	△	▲	△
Northern red oak	+	•	▲	△	▲	△
Northern white-cedar	•	•	▼	▼	▼	▼
Osage-orange	+	-	★		★	
Paper birch	•	•	●	○	▼	▼
Pignut hickory	•	-	▲	○	▲	△
Pin cherry*	•	-	▼	▼	▼	▼
Pitch pine	•	-	●	▼	●	▼
Post oak	+	-	▲	△	▲	△
Quaking aspen	•	•	▲	△	▲	△
Red maple	+	+	●	△	●	△
Red pine	-	-	●	▼	●	▼
Red spruce	-	+	▼	○	▼	○
Scarlet oak	•	•	●	○	▲	△
Serviceberry*	•	-	●	▼	●	▼
Shagbark hickory	•	-	▲	○	▲	○
Shortleaf pine	•	-	★		★	
Silver maple*	+	-	●	○	●	○
Southern red oak	+	-	★		★	
Striped maple	•	•	▼	▼	▼	▼
Sugar maple	+	+	●	△	●	△
Swamp white oak*	•	-	●	▼	▲	○
Sweet birch	-	•	▲	○	▲	○
Sweetbay	•	-	★		★	
Sweetgum	•	-	▲	△	▲	△
Sycamore*	•	-	▲	△	▲	△
Tamarack (native)	-	•	▼	▼	●	▼
Virginia pine	•	-	★		★	
White ash	-	•	▲	○	▲	○
White oak	+	•	▲	△	▲	△
White spruce	•	•	●	○	●	○
Willow oak*	•	-	★		★	
Yellow birch	•	+	▼	○	▼	○
Yellow-poplar	+	-	▲	△	▲	△

*Species with low model reliability based on five statistical metrics of the habitat models that affect change class. See maps and tables for more information (www.fs.fed.us/nrs/atlas/combined/resources/summaries).